

Jan Kurek, Włodzimierz M. Mikulski

Higher order jet prolongations type gauge natural bundles over vector bundles

Dedicated to Professor Andrzej Zajtz on the occasion of his 70th birthday with respect and gratitude

Abstract. Let $r \geq 3$ and $m \geq 2$ be natural numbers and E be a vector bundle with m -dimensional basis. We find all gauge natural bundles “similar” to the r -jet prolongation bundle $J^r E$ of E . We also find all gauge natural bundles “similar” to the vector r -tangent bundle $(J_{fi}^r(E, \mathbb{R})_0)^*$ of E .

Introduction

Natural bundles over m -manifolds were introduced by Nijenhuis [31] as a modern approach to geometric objects [1]. The most important results in the theory of natural bundles over m -manifolds is the Palais-Terng finite order theorem [32] and the Epstein-Thurston regularity theorem [7]. The sharp estimation of the order of natural bundles over m -manifolds was obtained by Zajtz [38]. From the order theorem follows that natural bundles over m -manifolds are associated to the principal bundles of repers of higher orders.

Natural bundles on some other categories over manifolds (e.g. all manifolds, fibered manifolds, principal fiber bundles, vector bundles, manifolds with structures) were introduced in [17]. Some sharp order estimations for some such natural bundles can be found in [17] and [22]. A natural bundle over all manifolds of infinite order can be found in [24]. Classifications of some type natural bundles over some categories over manifolds can be found in [35], [37], [3], [12], [23], [25], [26], [30], [16], [4], [6], [27], [18], [28].

Natural bundles make possible to precise define geometric constructions. Investigations, applications and classifications of natural operators (canonical constructions) on sections of natural bundles (geometric objects) were (and actually are) fundamental in the differential geometry. There are over 200 papers on this subject, e.g. [2], [3], [8]-[11], [13]-[15] [19]-[21], [33], [34], [36] and the fundamental monograph of Kolář, Michor and Slovák [17].

The present paper is a next contribution to the theory of natural bundles. Let us recall the following definition (see for ex. [17], [27]).

Let $F: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ be a covariant functor from the category \mathcal{VB}_m of vector bundles with m -dimensional bases and their vector bundle maps with local diffeomorphisms as base maps. Let $B_{\mathcal{VB}_m}: \mathcal{VB}_m \rightarrow \mathcal{M}f_m$ be the base functor.

A *gauge natural bundle over \mathcal{VB}_m* (called also a gauge bundle functor on \mathcal{VB}_m) is a functor F as above satisfying:

- (i) (**Base preservation**) $B_{\mathcal{VB}_m} \circ F = B_{\mathcal{VB}_m}$. Hence the induced projections form a functor transformation $\pi: F \rightarrow B_{\mathcal{VB}_m}$.
- (ii) (**Localization**) For every inclusion of an open vector subbundle $i_{E|U}: E|U \rightarrow E$, $F(E|U)$ is the restriction $\pi^{-1}(U)$ of $\pi: FE \rightarrow B_{\mathcal{VB}_m}(E)$ over U and $F i_{E|U}$ is the inclusion $\pi^{-1}(U) \rightarrow FE$.
- (iii) (**Regularity**) F transforms smoothly parametrized systems of \mathcal{VB}_m -morphisms into smoothly parametrized systems of \mathcal{VB}_m -morphisms.

A gauge natural bundle F over \mathcal{VB}_m is *fiber product preserving* if for every fiber product projections $E_1 \xleftarrow{pr_1} E_1 \times_M E_2 \xrightarrow{pr_2} E_2$ in the category \mathcal{VB}_m $FE_1 \xleftarrow{Fpr_1} F(E_1 \times_M E_2) \xrightarrow{Fpr_2} FE_2$ are fiber product projections in the category \mathcal{VB}_m . In other words $F(E_1 \times_M E_2) = F(E_1) \times_M F(E_2)$ modulo the corestriction of (Fpr_1, Fpr_2) .

All fiber product preserving gauge natural bundles over \mathcal{VB}_m are described in [28]. The most important example of the ones is the r -jet prolongation gauge natural bundle $J^r: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$. Another example is the (described in I.1) vector (r)-tangent gauge natural bundle $T^{(r)fl}: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$. Fiber product preserving gauge natural bundles on \mathcal{VB}_m with “similar to $T^{(r)fl}$ and J^r constructions” (Definitions 1 and 2) are also called vector r -tangent gauge natural bundles and r -jet prolongation gauge natural bundles over \mathcal{VB}_m , respectively.

Roughly speaking, the main results are the following classification theorems.

THEOREM A

Let $m \geq 2$ and $r \geq 3$ be integers. Up to isomorphism there are only two vector r -tangent gauge natural bundles over \mathcal{VB}_m .

THEOREM B

Let $m \geq 2$ and $r \geq 3$ be integers. Up to isomorphism there are only three r -jet prolongation gauge natural bundles over \mathcal{VB}_m .

The above theorems will be detailed formulated later (Theorems 1 and 2 of the present paper). In particular, the two vector r -tangent gauge natural

bundles over \mathcal{VB}_m and the three r -jet prolongation gauge natural bundles over \mathcal{VB}_m will be precisely constructed.

All manifolds and maps are assumed to be of class C^∞ . Manifolds are assumed to be finite dimensional.

1. The vector r -tangent gauge natural bundles over \mathcal{VB}_m

1. The vector (r) -tangent gauge natural bundle over \mathcal{VB}_m

Given a \mathcal{VB}_m -object $p: E \longrightarrow M$, the vector (r) -tangent bundle $T^{(r)fl}E$ of E is the vector bundle

$$T^{(r)fl}E = (J_{fl}^r(E, \mathbb{R})_0)^*$$

over M , where

$$J_{fl}^r(E, \mathbb{R})_0 = \{j_x^r \gamma \mid \gamma: E \longrightarrow \mathbb{R} \text{ is fiber linear, } \gamma_x = 0, x \in M\}.$$

Every \mathcal{VB}_m -map $f: E_1 \longrightarrow E_2$ covering $\underline{f}: M_1 \longrightarrow M_2$ induces a vector bundle map $T^{(r)fl}f: T^{(r)fl}E_1 \longrightarrow T^{(r)fl}E_2$ covering \underline{f} such that

$$\langle T^{(r)fl}f(\omega), j_{\underline{f}(x)}^r \xi \rangle = \langle \omega, j_x^r(\xi \circ f) \rangle,$$

$$\omega \in T_x^{(r)fl}E_1, j_{\underline{f}(x)}^r \xi \in J_{fl}^r(E_2, \mathbb{R})_0, x \in M_1.$$

The correspondence $T^{(r)fl}: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$ is a fiber product preserving gauge natural bundle.

2. Another construction of $T^{(r)fl}$

Let $p: E \longrightarrow M$ be a \mathcal{VB}_m -object. For any m -manifold M and $x \in M$ we have a canonical unital associative algebra homomorphism $t_x^{(r)}: J_x^r(M, \mathbb{R}) \longrightarrow \text{gl}((J_x^r(M, \mathbb{R})_0)^*)$ given by

$$t_x^{(r)}(j_x^r \gamma)(\omega)(j_x^r \eta) = \omega(j_x^r(\gamma \eta)),$$

$j_x^r \eta \in J_x^r(M, \mathbb{R})_0, j_x^r \gamma \in J_x^r(M, \mathbb{R}), \omega \in (J_x^r(M, \mathbb{R})_0)^*$. We have a vector bundle

$$\tilde{T}^{(r)fl}E = \bigcup_{x \in M} \text{Hom}_{t_x^{(r)}}(J^r \mathcal{C}_x^{\infty, fl}(E), (J_x^r(M, \mathbb{R})_0)^*)$$

over M . Here $\text{Hom}_{t_x^{(r)}}(J^r \mathcal{C}_x^{\infty, fl}(E), (J_x^r(M, \mathbb{R})_0)^*)$ is the vector space of all module homomorphisms over $t_x^{(r)}: J_x^r(M, \mathbb{R}) \longrightarrow \text{gl}((J_x^r(M, \mathbb{R})_0)^*)$ from the (free) $J_x^r(M, \mathbb{R})$ -module $J^r \mathcal{C}_x^{\infty, fl}(E)$ of r -jets at x of germs at x of fiber linear maps $E \longrightarrow \mathbb{R}$ into the $\text{gl}((J_x^r(M, \mathbb{R})_0)^*)$ -module $(J_x^r(M, \mathbb{R})_0)^*$. Every

\mathcal{VB}_m -map $f: E_1 \rightarrow E_2$ covering $\underline{f}: M_1 \rightarrow M_2$ induces a vector bundle map $\tilde{T}^{(r)fl} f: \tilde{T}^{(r)fl} E_1 \rightarrow \tilde{T}^{(r)fl} E_2$ covering \underline{f} such that

$$\tilde{T}^{(r)fl} f(\Phi)(j_{\underline{f}(x)}^r \xi)(j_{\underline{f}(x)}^r \gamma) = \Phi(j_x^r(\xi \circ f))(j_x^r(\gamma \circ \underline{f})),$$

$\Phi \in \text{Hom}_{t_x^{(r)}}(J^r C_x^{\infty, fl}(E_1), (J_x^r(M_1, \mathbb{R})_0)^*)$, $x \in M_1$, $j_{\underline{f}(x)}^r \xi \in J^r C_{\underline{f}(x)}^{\infty, fl}(E_2)$, $j_{\underline{f}(x)}^r \gamma \in J^r(M_2, \mathbb{R})_0$.

The correspondence $\tilde{T}^{(r)fl}: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ is a fiber product preserving gauge natural bundle.

LEMMA 1

We have a natural isomorphism $T^{(r)fl} \simeq \tilde{T}^{(r)fl}$

Proof. Define a base preserving vector bundle map $\Theta: T^{(r)fl} E \rightarrow \tilde{T}^{(r)fl} E$ by

$$\Theta(\omega)(j_x^r \xi)(j_x^r \gamma) = T^{(r)fl}(\xi)(\omega)(j_x^r \gamma) \in \mathbb{R},$$

$\omega \in T_x^{(r)fl} E$, $j_x^r \xi \in J^r C_x^{\infty, fl}(E)$, $j_x^r \gamma \in J_x^r(M, \mathbb{R})_0$, $x \in M$, where $\xi: E \rightarrow \mathbb{R}$ is considered as $\mathcal{VB}_{m,n}$ -map $\xi: E \rightarrow M \times \mathbb{R}$, $\xi(v) = (p(v), \xi(v))$, $v \in E$ and where $\gamma: M \rightarrow \mathbb{R}$ is considered as the fiber linear map $\gamma: M \times \mathbb{R} \rightarrow \mathbb{R}$, $\gamma(m, t) = \gamma(m)t$, $m \in M$, $t \in \mathbb{R}$. Using trivializations of E it is easy to see that Θ is a vector bundle isomorphism.

3. The vector $[r]$ -tangent gauge natural bundle over \mathcal{VB}_m

Given a \mathcal{VB}_m -object $p: E \rightarrow M$ we have the vector bundle

$$T^{[r]fl} E = E \otimes (J^r(M, \mathbb{R})_0)^*$$

over M . Every \mathcal{VB}_m -map $f: E_1 \rightarrow E_2$ covering $\underline{f}: M_1 \rightarrow M_2$ induces (in obvious way) a vector bundle map $T^{[r]fl} f: T^{[r]fl} E_1 \rightarrow T^{[r]fl} E_2$ covering \underline{f} .

The correspondence $T^{[r]fl}: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ is a fiber product preserving gauge natural bundle.

4. Another construction of $T^{[r]fl}$

For any m -manifold M and $x \in M$ we have a canonical unital associative algebra homomorphism $t_x^{[r]}: J_x^r(M, \mathbb{R}) \rightarrow \text{gl}((J_x^r(M, \mathbb{R})_0)^*)$ given by

$$t_x^{[r]}(j_x^r \gamma)(\omega) = \gamma(x)\omega,$$

$j_x^r \gamma \in J_x^r(M, \mathbb{R})$, $\omega \in (J_x^r(M, \mathbb{R})_0)^*$. Using $t_x^{[r]}$ instead of $t_x^{(r)}$ we can construct similarly as in Section 2 a fiber product preserving gauge natural bundle $\tilde{T}^{[r]fl}: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$.

LEMMA 2

We have a natural isomorphism $T^{[r]fl} \cong \tilde{T}^{[r]fl}$.

Proof. The proof is quite similar to the one of Lemma 1.

5. $T^{(r)fl} \neq T^{[r]fl}$

LEMMA 3

The gauge natural bundle $T^{(r)fl}: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ and $T^{[r]fl}: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ are not isomorphic.

Proof. Let $f: \mathbb{R}^{m,n} \rightarrow \mathbb{R}^{m,n}$ be a $\mathcal{VB}_{m,n}$ -map given by

$$f(x, y) = (x, y + x^1 y), \quad x = (x^1, \dots, x^m) \in \mathbb{R}^m, \quad y \in \mathbb{R}^n.$$

Clearly $T_0^{[r]fl} f = \text{id}$ and $T_0^{(r)fl} f \neq \text{id}$. That is why there is no base preserving vector bundle isomorphism $\Theta: T^{(r)fl} \mathbb{R}^{m,n} \rightarrow T^{[r]fl} \mathbb{R}^{m,n}$ commuting with f .

6. The vector r -tangent gauge natural bundles over \mathcal{VB}_m

DEFINITION 1

We say that a fiber product preserving gauge natural bundle $F: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ is a vector r -tangent gauge natural bundle if modulo isomorphism F can be constructed similarly as $\tilde{T}^{(r)fl}$ by using eventually another canonical unital associative algebra homomorphism $t_x: J_x^r(M, \mathbb{R}) \rightarrow \text{gl}((J_x^r(M, \mathbb{R})_0)^*)$ instead of $t_x^{(r)}$.

We have the following classification theorem.

THEOREM 1

Let $m \geq 2$ and $r \geq 3$ be integers. Up to isomorphism $T^{(r)fl}: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ and $T^{[r]fl}: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ are only vector r -tangent gauge natural bundles over \mathcal{VB}_m .

Proof. Suppose that for any m -manifold M and $x \in M$

$$t_x: J_x^r(M, \mathbb{R}) \rightarrow \text{gl}((J_x^r(M, \mathbb{R})_0)^*)$$

is a canonical unital associative algebra homomorphism. By the invariance of t_0 with respect to $(\tau^1 x^1, \dots, \tau^m x^m)$ for $\tau^i \neq 0$ we deduce that

$$t_0(j_0^r x^1)((j_0^r x^\alpha)^*)(j_0^r x^2) = 0$$

for $\alpha \neq (1, 1, 0, \dots, 0)$. Then we can write

$$t_0(j_0^r x^1)(\omega)(j_0^r x^2) = a\omega(j_0^r(x^1 x^2))$$

for all $\omega \in (J_0^r(\mathbb{R}^m, \mathbb{R})_0)^*$, where $a = t_0(j_0^r x^1)((j_0^r(x^1 x^2))^*)(j_0^r x^2)$. Then by the invariance of t_0 with respect to 0-preserving diffeomorphisms we have

$$t_0(j_0^r \gamma)(\omega)(j_0^r \eta) = a\omega(j_0^r(\gamma\eta))$$

for all $\omega \in (J_0^r(\mathbb{R}^m, \mathbb{R})_0)^*$ and all $j_0^r \gamma, j_0^r \eta \in J_0^r(\mathbb{R}^m, \mathbb{R})_0$. But t_0 is an algebra homomorphism. Then

$$\begin{aligned} a\omega(j_0^r((x^1)^2 x^2)) &= t_0(j_0^r((x^1)^2))(\omega)(j_0^r x^2) \\ &= t_0(j_0^r x^1)(t_0(j_0^r x^1)(\omega))(j_0^r x^2) \\ &= at_0(j_0^r x^1)(\omega)(j_0^r(x^1 x^2)) \\ &= a^2\omega(j_0^r((x^1)^2 x^2)). \end{aligned}$$

If $r \geq 3$ then $a^2 = a$, i.e. $a = 1$ or $a = 0$. Hence $t_x = t_x^{(r)}$ or $t_x = t_x^{[r]}$.

II. The r -jet prolongation gauge natural bundles over \mathcal{VB}_m

7. The (usual) r -jet prolongation gauge natural bundle J^r over \mathcal{VB}_m

Given a \mathcal{VB}_m -object $p: E \rightarrow M$ the (usual) r -jet prolongation $J^r E$ of E is a vector bundle

$$J^r E = \{j_x^r \sigma \mid \sigma \text{ is a local section of } E, x \in M\}$$

over M . Every \mathcal{VB}_m -map $f: E_1 \rightarrow E_2$ covering $\underline{f}: M_1 \rightarrow M_2$ induces a vector bundle map $J^r f: J^r E_1 \rightarrow J^r E_2$ covering \underline{f} such that

$$J^r f(j_x^r \sigma) = j_{\underline{f}(x)}^r(f \circ \sigma \circ \underline{f}^{-1}), \quad j_x^r \sigma \in J^r E_1.$$

The functor $J^r: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ is a fiber product preserving gauge natural bundle.

8. Another construction of J^r

Let $p: E \rightarrow M$ be a \mathcal{VB}_m -object. For any m -manifold M and $x \in M$ we have a canonical unital associative algebra homomorphism $t_x^{r,m}: J_x^r(M, \mathbb{R}) \rightarrow \text{gl}(J_x^r(M, \mathbb{R}))$ given by

$$t_x^{r,m}(j_x^r \eta)(j_x^r \gamma) = j_x^r(\gamma\eta),$$

$j_x^r \eta \in J_x^r(M, \mathbb{R}), j_x^r \gamma \in J_x^r(M, \mathbb{R})$. We have a vector bundle

$$\tilde{J}^r E = \bigcup_{x \in M} \text{Hom}_{t_x^{r,m}}(J^r \mathcal{C}_x^{\infty, fl}(E), J_x^r(M, \mathbb{R}))$$

over M . Here $\text{Hom}_{t_x^{r,m}}(J^r \mathcal{C}_x^{\infty,fl}(E), J_x^r(M, \mathbb{R}))$ is the vector space of all module homomorphisms over $t_x^{r,m}: J_x^r(M, \mathbb{R}) \rightarrow \text{gl}(J_x^r(M, \mathbb{R}))$ from the (free) $J_x^r(M, \mathbb{R})$ -module $J^r \mathcal{C}_x^{\infty,fl}(E)$ of r -jets at x of germs at x of fiber linear maps $E \rightarrow \mathbb{R}$ into the $\text{gl}(J_x^r(M, \mathbb{R}))$ -module $J_x^r(M, \mathbb{R})$. Every \mathcal{VB}_m -map $f: E_1 \rightarrow E_2$ covering $\underline{f}: M_1 \rightarrow M_2$ induces a vector bundle map $\tilde{J}^r f: \tilde{J}^r E_1 \rightarrow \tilde{J}^r E_2$ covering \underline{f} such that

$$\tilde{J}^r f(\Phi)(j_{\underline{f}(x)}^r \xi) = J^r(\underline{f}, \text{id}_{\mathbb{R}}) \circ \Phi(j_x^r(\xi \circ f)),$$

$\Phi \in \text{Hom}_{t_x^{r,m}}(J^r \mathcal{C}_x^{\infty,fl}(E_1), J_x^r(M_1, \mathbb{R}))$, $x \in M_1$, $j_{\underline{f}(x)}^r \xi \in J^r \mathcal{C}_{\underline{f}(x)}^{\infty,fl}(E_2)$.

The correspondence $\tilde{J}^r: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ is a fiber product preserving gauge natural bundle.

LEMMA 4

We have a natural isomorphism $J^r \cong \tilde{J}^r$.

Proof. The proof is similar to the one of Lemma 1. We define an isomorphism $\Theta: J^r E \rightarrow \tilde{J}^r E$ by

$$\Theta(j_x^r \sigma)(j_x^r \xi) = j_x^r(\xi \circ \sigma),$$

$j_x^r \sigma \in J_x^r E$, $j_x^r \xi \in J^r \mathcal{C}_x^{\infty,fl}(E)$.

9. The vertical r -jet prolongation gauge natural bundle J_v^r over \mathcal{VB}_m

Given a \mathcal{VB}_m -object $p: E \rightarrow M$, the vertical r -jet prolongation $J_v^r E$ of E is a vector bundle

$$J_v^r E = \{j_x^r \sigma \mid \sigma: M \rightarrow E_x, x \in M\}$$

over M . Every \mathcal{VB}_m -map $f: E_1 \rightarrow E_2$ covering $\underline{f}: M_1 \rightarrow M_2$ induces a vector bundle map $J_v^r f: J_v^r E_1 \rightarrow J_v^r E_2$ covering \underline{f} such that

$$J_v^r f(j_x^r \sigma) = j_{\underline{f}(x)}^r(f \circ \sigma \circ \underline{f}^{-1}), \quad j_x^r \sigma \in J_v^r E_1.$$

The functor $J_v^r: \mathcal{VB}_m \rightarrow \mathcal{VB}_m$ is a fiber product preserving gauge natural bundle.

10. Another construction of J_v^r

For any m -manifold M and $x \in M$ we have a canonical unital associative algebra homomorphism $t_x^{r,m,o}: J_x^r(M, \mathbb{R}) \rightarrow \text{gl}(J_x^r(M, \mathbb{R}))$ given by

$$t_x^{r,m,o}(j_x^r \eta)(j_x^r \gamma) = \eta(x) j_x^r \gamma,$$

$j_x^r \gamma \in J_x^r(M, \mathbb{R})$, $j_x^r \eta \in J_x^r(M, \mathbb{R})$. Using $t_x^{r,m,o}$ instead of $t_x^{r,m}$ we can construct similarly as in Section 9 a fiber product preserving gauge natural bundle $\tilde{J}_v^r: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$.

LEMMA 5

We have a natural isomorphism $J_v^r \cong \tilde{J}_v^r$.

Proof. The proof is quite similar to the one of Lemma 4.

11. The $[r]$ -jet prolongation gauge natural bundle $J^{[r]}$ over \mathcal{VB}_m

For any m -manifold M and $x \in M$ we have a canonical unital associative algebra homomorphism $t_x^{[r],m}: J_x^r(M, \mathbb{R}) \longrightarrow \text{gl}(J_x^r(M, \mathbb{R}))$ given by

$$t_x^{[r],m}(j_x^r \eta)(j_x^r \gamma) = j_x^r(\eta\gamma) - \gamma(x)j_x^r \eta + \eta(x)\gamma(x)j_x^r 1,$$

$j_x^r \gamma \in J_x^r(M, \mathbb{R})$, $j_x^r \eta \in J_x^r(M, \mathbb{R})$. Using $t_x^{[r],m}$ instead of $t_x^{r,m}$ we can construct similarly as in Section 9 a fiber product preserving gauge natural bundle $J^{[r]}: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$.

12. J^r , J_v^r and $J^{[r]}$ are not equivalent

LEMMA 6

The gauge natural bundles $J^r: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$ and $J_v^r: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$ are not isomorphic.

Proof. Let $f: \mathbb{R}^{m,n} \longrightarrow \mathbb{R}^{m,n}$ be a $\mathcal{VB}_{m,n}$ -map given by

$$f(x, y) = (x, y + x^1 y), \quad x = (x^1, \dots, x^m) \in \mathbb{R}^m, \quad y \in \mathbb{R}^n.$$

Clearly $J_0^r f \neq \text{id}$ and $(J_v^r)_0 f = \text{id}$. That is why there is no base preserving vector bundle isomorphism $\Theta: J^r \mathbb{R}^{m,n} \longrightarrow J_v^r \mathbb{R}^{m,n}$ commuting with f .

LEMMA 7

The gauge natural bundles $J^{[r]}: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$ and $J_v^r: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$ are not isomorphic.

Proof. The proof is quite similar to the one of Lemma 6.

LEMMA 8

Let $m \geq 2$. The gauge natural bundles $J^{[r]}: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$ and $J^r: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$ are not isomorphic.

Proof. It follows from some result from [29]. More precisely, the vector space of all $\mathcal{VB}_{m,n}$ -natural affinors on $J^r E$ is one dimensional and the vector space of all $\mathcal{VB}_{m,n}$ -natural affinors on $J^{[r]} E$ is two dimensional.

13. The r -jet prolongation gauge natural bundles over \mathcal{VB}_m

DEFINITION 2

We say that a fiber product preserving gauge natural bundle $F: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$ is a r -jet prolongation gauge natural bundle if modulo isomorphism F can be constructed similarly as \tilde{J}^r by using eventually another canonical unital associative algebra homomorphism $t_x: J_x^r(M, \mathbb{R}) \longrightarrow \text{gl}(J_x^r(M, \mathbb{R}))$ instead of $t_x^{r,m}$.

We have the following classification theorem.

THEOREM 2

Let $m \geq 2$ and $r \geq 3$ be integers. Up to isomorphism $J^r: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$, $J_v^r: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$ and $J^{[r]}: \mathcal{VB}_m \longrightarrow \mathcal{VB}_m$ are only r -jet prolongation gauge natural bundles over \mathcal{VB}_m .

14. A preparation

LEMMA 9

For any $\alpha \in \mathbb{R}$ we have a canonical unital algebra homomorphism

$$t_x^{<\alpha>}: J_x^r(M, \mathbb{R}) \longrightarrow \text{gl}(J_x^r(M, \mathbb{R}))$$

given by

$$t_x^{<\alpha>}(j_x^r \gamma)(j_x^r \eta) = j_x^r(\gamma \eta) + \alpha \eta(x) j_x^r \gamma - \alpha \eta(x) \gamma(x) j_x^r 1,$$

where $j_x^r \eta, j_x^r \gamma \in J_x^r(M, \mathbb{R})$, $\dim(M) = m$, $x \in M$. If $\alpha \notin \{-1, 0\}$ we have a canonical isomorphism $a_x: J_x^r(M, \mathbb{R}) \longrightarrow J_x^r(M, \mathbb{R})$ given by

$$a_x(j_x^r \gamma) = -\frac{\alpha + 1}{\alpha} j_x^r \gamma + \gamma(x) j_x^r 1$$

such that $t_x^{<\alpha>}(j_x^r \eta) \circ a_x = a_x \circ t_x^{r,m}(j_x^r \eta)$ for any $j_x^r \eta \in J_x^r(M, \mathbb{R})$. Roughly speaking, for $\alpha \notin \{-1, 0\}$ homomorphism $t_x^{<\alpha>}$ is canonically isomorphic to $t_x^{r,m}$.

Proof. It is easy to verify.

15. Proof of Theorem 2

Let for any m -manifold M and $x \in M$

$$t_x: J_x^r(M, \mathbb{R}) \longrightarrow \text{gl}(J_x^r(M, \mathbb{R}))$$

be a canonical unital associative algebra homomorphism. Clearly, t_x is uniquely determined by the values

$$t_0(j_0^r \gamma)(j_0^r \eta) \in J_0^r(\mathbb{R}^m, \mathbb{R})$$

for any $j_0^r \gamma, j_0^r \eta \in J_0^r(\mathbb{R}^m, \mathbb{R})$ with $\gamma(0) = 0$. By the invariance of t_0 with respect to 0-preserving diffeomorphisms and the rank theorem and $m \geq 2$ we can assume that

$$j_0^r \gamma = j_0^r x^1$$

and

$$j_0^r \eta = j_0^r 1 \quad \text{or} \quad j_0^r \eta = j_0^r x^2.$$

By the invariance of t_0 with respect to $(\tau_1 x^1, \dots, \tau_m x^m)$ for $\tau_1, \dots, \tau_m \in \mathbb{R}_+$ we deduce that

$$t_0(j_0^r x^1)(j_0^r x^2) = \sigma j_0^r(x^1 x^2)$$

and

$$t(j_0^r x^1)(j_0^r 1) = \rho j_0^r x^1$$

for some $\sigma, \rho \in \mathbb{R}$. Then by the invariance of t_0 with respect to 0-preserving diffeomorphisms we have

$$t_0(j_0^r \xi)(j_0^r \eta) = \sigma j_0^r(\xi \eta)$$

and

$$t_0(j_0^r \xi)(j_0^r 1) = \rho j_0^r \xi$$

for all $j_0^r \xi \in J_0^r(\mathbb{R}^m, \mathbb{R})$ and $j_0^r \eta \in J_0^r(\mathbb{R}^m, \mathbb{R})$ and with $\xi(0) = \eta(0) = 0$. Then, since t_0 is an unital homomorphism, it is easy to verify (using the assumptions on r, m) that $\sigma = \rho = 0$ or $\sigma = 1$ and ρ is arbitrary. Therefore $t_x = t_x^{r,m,\sigma}$ or $t_x = t_x^{<\alpha>}$ for $\alpha = \rho - 1$.

Using Lemma 9 we end the proof.

References

- [1] J. Aczél, S. Gołąb, *Funktionalgleichungen der Theorie der geometrischen Objekte*, Monografie Matematyczne, Warszawa, PWN, 1960.
- [2] J. Dębecki, J. Gancarzewicz, M. de Leon, W.M. Mikulski, *Invariants of Lagrangians and their classifications*, J. Math. Phys. **35**(9) (1994), 4568-4593.
- [3] M. Doupovec, *Natural operators transforming vector fields to the second order tangent bundle*, Časopis Pěst. Mat. **115** (1990), 64-72.
- [4] M. Doupovec, I. Kolář, *On the jets of fibered manifold morphisms*, Cahiers Topologie Géom. Différentielle Catég. **40** (1999), 21-30.
- [5] D.J. Eck, *Product-preserving functors on smooth manifolds*, J. Pure Appl. Algebra **42** (1986), 133-140.
- [6] D.J. Eck, *Gauge natural bundles and generalized gauge theories*, Mem. Amer. Math. Soc. **33** (1981), no. **247**.
- [7] D.B.A. Epstein, W.P. Thurston, *Transformation groups and natural bundles*, Proc. London Math. Soc. (3) **38** (1979), 219-236.

- [8] J. Gancarzewicz, *Liftings of functions and vector fields to natural bundles*, Dissertationes Math. **212** (1983).
- [9] J. Gancarzewicz, I. Kolář, *Some gauge-natural operators on linear connections*, Monatsh. Math. **111** (1991), 23-33.
- [10] J. Gancarzewicz, W. Mikulski, Z. Pogoda, *Lifts of some tensor fields and connections to a product preserving functor*, Nagoya Math. J. **135** (1994), 1-41.
- [11] J. Janyška, D. Krupka, *Lectures on differential invariants*, Folia Facultatis Scientiarum Naturalium Universitatis Purkynianae Brunensis. Mathematica, 1., 1990.
- [12] G. Kainz, P. Michor, *Natural transformations in differential geometry*, Czechoslovak Math. J. **37(112)** (1987), 584-607.
- [13] I. Kolář, *On natural operators on vector fields*, Ann. Global Anal. Geom. **6(2)** (1988), 109-117.
- [14] I. Kolář, *Gauge-natural forms of Chern-Weil type*, Ann. Global Anal. Geom. **11** (1993), 41-47.
- [15] I. Kolář, *Natural operators related with the variational calculus*, Differential Geom. and Its Appl. Proceedings Opava (1993), 461-472.
- [16] I. Kolář, *Bundle functors of jet type*, Differential Geom. and Appl. Proc. of the 7th International Conference Brno (1998), 231-237.
- [17] I. Kolář, P.W. Michor, J. Slovák, *Natural Operations in Differential Geometry*, Springer Verlag, Berlin, 1993.
- [18] I. Kolář, W.M. Mikulski, *On the fiber product preserving bundle functors*, Differential Geom. Appl. **11** (1999), 105-115.
- [19] J. Kurek, *On the natural operators of Bianchi type*, Arch. Math. (Brno) **27a** (1991), 25-29.
- [20] J. Kurek, W.M. Mikulski, *The natural operators lifting 1-forms to some vector bundle functors*, Colloq. Math. **93(2)** (2002), 259-265.
- [21] M. Kureš, W.M. Mikulski, *Natural operators lifting vector fields to bundles of Weil contact elements*, Czechoslovak Math. J. **54(129)** (2004), no. 4, 855-867.
- [22] K. Lubaš, A. Zajtz, *Estimates of the action order of some jet groups*, Proc. Conf. Differential Geometry and Appl. Brno (1998), Masaryk University in Brno (1999), 247-257.
- [23] O.O. Luciano, *Categories of multiplicative functors and Morimoto's conjecture*, Preprint 46, Inst. Fourier, Laboratoire des Mathématiques, Grenoble, 1986.
- [24] W.M. Mikulski, *There exists a prolongation functor of infinite order*, Časopis Pěst. Mat. **114** (1989), 57-59.
- [25] W.M. Mikulski, *Product preserving bundle functors on fibered manifolds*, Arch. Math. (Brno) **32(4)** (1996), 307-316.
- [26] W.M. Mikulski, *On the product preserving bundle functors on k -fibered manifolds*, Demonstratio Math. **34(3)** (2001), 693-700.
- [27] W.M. Mikulski, *Product preserving gauge bundle functors on vector bundles*, Colloq. Math. **90(2)** (2001), 277-285.

- [28] W.M. Mikulski, *On the fiber product preserving gauge bundle functors on vector bundles*, Ann. Polon. Math. **82** (2003), no. 3, 251-264.
- [29] W.M. Mikulski, *The natural affinors on the r -jet prolongations of a vector bundle*, Demonstratio Math. **37** (2004), no. 3, 709-717.
- [30] W.M. Mikulski, J. Tomáš, *Product preserving bundle functors on fibered manifolds*, Colloq. Math. **96** (2003), no. 1, 17-26.
- [31] A. Nijenhuis, *Natural bundles and their general properties*, Geometric objects revisited. Differential geometry (in honor of Kentaro Yano), pp. 317-334. Kinokuniya, Tokyo, 1972.
- [32] R.S. Palais, C.-L. Terng, *Natural bundles have finite order*, Topology **16** (1977), 271-277.
- [33] M. Paluszny, A. Zajtz, *Foundation of the geometry of natural bundles*, Lect. Notes Univ. Caracas, 1984.
- [34] J. Slovák, *Peetre theorem for nonlinear operators*, Ann. Global Anal. Geom. **6** (1988), 273-283.
- [35] A. Zajtz, *Geometric object with finitely determined germs*, Ann. Polon. Math. **49** (1988), 157-168.
- [36] A. Zajtz, *On the order of natural differential operators and liftings*, Ann. Polon. Math. **49** (1988), 169-178.
- [37] A. Zajtz, *A classification of natural bundles*, Colloq. Math. Soc. János Bolyai, **46**, Topics in differential geometry, Vol. I, II (Debrecen, 1984), 1333-1357.
- [38] A. Zajtz, *The sharp upper bound on the order of natural bundles of given dimension*, Bull. Soc. Math. Belg. Sér. B **39** (1987), 347-357.

*Institute of Mathematics
Maria Curie Skłodowska University
Plac Marii Curie Skłodowskiej 1
PL-20-031 Lublin
Poland
e-mail: kurek@golem.umcs.lublin.pl*

*Institute of Mathematics
Jagiellonian University
Reymonta 4
PL-30-059 Kraków
Poland
e-mail: mikulski@im.uj.edu.pl*